

Annual Conference on

MICROBIAL PATHOGENESIS, INFECTIOUS DISEASE, ANTIMICROBIALS AND DRUG RESISTANCE

August 23-24, 2017 | Toronto, Canada

Phytochemical screening of the exudate of *Aloe otallensis* and its effect on *Leishmania donovani* parasite

Tesfaye Zerihun
Addis Ababa University, Ethiopia

Objectives: The objectives of the study is to evaluate antileishmanial activity of methanolic extract of *Aloe otallensis* (*A. otallensis*) on the promastigote stage of *Leishmania donovani* (*L. donovani*) as compared to standard drugs and to screen its phytochemical constituents.

Methods: Phytochemical screening was done by using the method mentioned by Evans and Trease on methanolic extract of the exudates of *Aloe otallensis* leaves. The extract was also evaluated for in vitro antileishmanial activity against *L. donovani* which is found from the Parasitology Unit of Black Lion Hospital. The result was compared to standard drugs of sodium stibogluconate, milfostin and paramomycin.

Results: The extract has a good antileishmanial activity with an IC₅₀ of 0.123 0 µg/mL on *L. donovani* (AM 563).

The experimental data showed that relatively it had better activity than paramomycin and milfostin but less activity than sodium stibogluconate. The data analyses were done by GraphPad Prism version 5 software after it was read by ELISA reader at the wave length of 650 nm. The phytochemical screening of the exudates of *A. otallensis* showed the presence of phenol, alkaloid and saponin.

Conclusions: The methanol extract of the exudates of *A. otallensis* has a good anti-leishmaniasis activity and this may be attributed to phenol, alkaloid and saponin present in the plant. But it needs further analysis for the conformation of which constituent presents in high concentration to know which one has the strongest effect.

e: zerihun.tesfaye@aau.edu.et